

INDEX TO VOLUME I.

- ACANTHIS**, 145.
 canescens, 152.
 canescens exilipes, 152.
 hornemannii, 152.
 hornemannii exilipes, 152.
 linaria, 152.
 linaria cabaret, 153.
 linaria holboellii, 153.
 linaria pallescens, 153.
Accipiter, 58.
 cooperi, 247.
 fuscus, 247, 295.
Acredula caudata, 342.
Acrocephalus dybowskii, 82.
Actiturus, 142.
 bartramius, 315.
Actodromas acuminata, 78.
 maculatus, 218.
Adney, E. T., the Cardinal Grosbeak breeding in Brooklyn, N. Y., 390.
Æchmophorus major, 316.
Ægialites albidipectus, 83.
 melodus circumcinctus, 397.
 mongolicus, 78.
Ægialitis collaris, 313.
 falklandica, 313.
Ægiothus, 145, 146.
 canescens canescens, 152.
 canescens exilipes, 152.
 exilipes, 152, 159.
 fuscescens, 152.
 hornemannii, 152.
 linarius, 152, 159.
 linarius exilipes, 152.
 linarius hoboellii, 153.
 rostratus, 153.
 rufescens, 152.
Agelæus phœniceus, 245.
Aguila, 109, 110.
 rojizo, 109.
Aix, 46, 144.
 sponsa, 187, 249.
Alauda ludoviciana, 168.
 minor, 260.
 mongolica, 228.
 pensylvanicus, 167, 168.
 rubra, 168.
 rufa, 261.
Albicilla, 140.
Alcon, 30.
Alconcito, 110.
Alle nigricans, 35, 115.
Allen, J. A., Zoölogical Nomenclature, 338.
Alouette aux joues brunes de Pensylvanie, 167, 168.
Aluco flammeus, 29.
 flammeus americanus, 246.
Aluette de Pensylvanie, 167, 168.
American Ornithologists' Union, Second Meeting of, 369.
Ammodromus maritimus, 176.
Ampelis cedrorum, 176, 244.
 garrula, 158, 244.
Anas boscas, 248.
 boschas, 122.
 obscura, 248.
Anorthura hiemalis, 15.
 pallescens, 82.
 troglodytes hiemalis, 138.
Anotus, 144.
Anser, 143.
 albifrons gambeli, 248.
Anthus japonicus, 82.
 ludovicianus, 89, 121, 176, 186.
 pensilvanicus, 168, 209.
 pratensis, 77.
 stejnegeri, 82.
Antrostomus parvulus, 24.
Apteryx australis, 81.
Aquila, 140.
 chrysaëtus canadensis, 79, 247.
Aramides ypecaha, 276.
Aramis pictus, 115.
 scolopaceus, 277.
Archibuteo lagopus sancti-johannis, 79, 247, 397.
Ardea cocoi, 271.
 candidissima, 271.
 egretta, 271.
 herodias, 247.
 occidentalis, 162.
 sibilatrix, 271.
 wardi, 161.
 würdemanni, 162.
Ardetta exilis, 248.

- Ardetta involucris*, 271.
Arenaria interpres, 229.
 melanocephala, 229.
Arquatella, 142.
 couesi, 77.
 maritima, 76.
Asio accipitrinus, 29, 115.
 brachyotus, 115.
Astrigalinus tristis, 244, 328.
Astur atricapillus, 247.
 atricapillus henshawi, 166.
 atricapillus striatulus, 252.
 badius, 340.
 brevipes, 340.
 poliopsis, 340.
 polyzonoides, 340.
 sphenurus, 340.
Asturina pucherani, 30.
Atthis, 57.
Auk, Black-throated, 98.
 Great, 81, 208.
Auriparus flaviceps, 354.
Avestruz, 318.
Aythya americana, 249.
 BAILEY, H. B., breeding habits of
 the Everglade Kite, 95.
 Baird, Brewer, and Ridgway, notice
 of their work on the Water Birds
 of North America, 382.
 Banks, J. W., nesting of the Broad-
 winged Hawk (*Buteo pennsylvanicus*), 95; the Great Horned
 Owl (*Bubo virginianus*) in con-
 finement, 194.
 Barrows, W. B., birds of the Low-
 er Uruguay, 20, 109, 270, 313;
 abnormal coloration in a caged
 Robin, 90.
 Batchelder, C. F., description of
 the first plumage of Clarke's
 Crow, 16; Buffon's Skua in Ver-
 mont, 97.
 Becasino, 314.
 Belding, L., notices of his papers on
 birds of Lower California, etc., 83,
 284.
 Bernicla, 143.
 canadensis, 80, 248, 309.
 canadensis hutchinsi, 80.
 canadensis leucoparia, 80.
 canadensis occidentalis, 80.
 brenta, 248.
 brenta nigricans, 96.
 nigricans, 187.
 Bicknell, E. P., a Study of the Sing-
 ing of our birds, 60, 126, 209, 322.
 Bird, Cardinal, 189.
 Cedar, 6, 294.
 Chipping, 190.
 Bird, Cow. See Cowbird.
 Guachera, 307.
 Indigo, 6, 67, 86.
 Phoebe, 65.
 Summer Red, 290.
 Thistle, 294.
 Bittern, American, 247.
 Least, 188, 248.
 Blackbird, Cow. See Cowbird.
 Crow, 6, 86, 185.
 Red-winged, 6, 245.
 Yellow-headed, 245, 293.
 Blackhead, Ring-billed, 249.
 Bluebird, 6, 13, 86, 133, 244.
 Californian, 121.
 Bobolink, 6, 8, 64, 245.
Bolborhynchus monachus, 28.
Bonasa umbellus, 186, 247.
Botaurus, 143.
 lentiginosus, 87, 247.
 mugitans, 115.
Brachyrhampus brachypterus, 386.
 hypoleucus, 396.
 kittlitzii, 81.
 Brant, 248.
 Black, 187.
 Brewer, T. M., see Baird, Brewer,
 and Ridgway.
 Brewster, W., notes on the Sum-
 mer Birds of Berkshire County,
 Mass., 5; notice of his annota-
 tions to the 'Birds and Feathered
 Fowles' of Morton's 'New English
 Canaan,' 84; on a specimen of
 Helminthophaga leucobronchialis
 from Connecticut, 91; recent oc-
 currence of the Black-backed
 Three-toed Woodpecker in Mas-
 sachusetts, 93; a singular speci-
 men of the Black-and-White
 Creeper, 190; note on the Lesser
 Glaucous-winged Gull in New
 York, 196; an ornithological
 swindler, 295; notice of his Birds
 observed in the Gulf of St. Law-
 rence, 379.
 Brown, N. C., a second season in
 Texas, 120; the Widgeon in Maine
 in February, 394.
 Browne, F. C., the Loggerhead
 Shrike again in Massachusetts,
 291; *Geothlypis trichas* wintering
 in Massachusetts, 389.
 Bryant, W. E., Nest and eggs of
 Myiadestes townsendi, 91.
 Bubo, 57.
 virginianus, 29, 194, 246.
 virginianus subarcticus, 79.
 Budytes flava, 77.

- Buffalo Naturalists' Field Club,
 notice of its 'Bulletin,' 184.
 Buho, 29.
 Bunting, Cow, 189.
 Lark, 286, 293.
 Snow, 159, 245, 294.
 Towhee, 13, 245.
 Buteo, 140.
 abbreviatus, 122.
 borealis, 123, 247.
 borealis krideri, 100, 123.
 cooperi, 253.
 erythronotus, 109.
 harlani, 253.
 pennsylvanicus, 95, 188.
 pteroles, 109.
 swainsoni, 247.
 Butorides cyanurus, 271.
 Butterball, 249.
 Buzzard, Turkey, 187, 247, 286, 293.
 CABURÉ, 29.
 Calamospiza bicolor, 121, 293.
 Calcarius, 118.
 nivalis, 118.
 Calendula, 56.
 Calidris, 142.
 arenaria, 397.
 Calodromas elegans, 318.
 Calypste costæ, 354.
 Calyptophilus, 3.
 frugivorus, 3.
 Campephilus boiaei, 25.
 Campotolæmus labradorius, 207.
 Canace canadensis, 15, 247.
 Caprimulgus vociferus, 246.
 Cardinal, Texas, 122.
 Cardinalis ruber, 172.
 ruber carneus, 172.
 ruber coccineus, 172.
 ruber igneus, 172.
 virginianus, 172, 390.
 Carpenter, J. H., the nest of the
 Saw-whet Owl, 94.
 Carpinterito, 25.
 Carpintero, 20, 25.
 blanco, 25.
 Carpodacus cassini, 288.
 frontalis, 288, 289, 355.
 purpureus, 288, 327.
 Carrancho, 111.
 Catbird, 6, 13, 132, 185, 244.
 Cathartes atratus, 113, 285.
 aura, 87, 113, 247, 293.
 Chætura pelagica, 161, 246.
 Chaja, 273.
 Certhia familiaris, 281.
 familiaris β americana, 281.
 familiaris rufa, 121, 136, 157.
 Ceryle alcyon, 179, 246.
 amazona, 26.
 americana, 26.
 torquata, 26.
 Chamberlain, M., nest and eggs of
 the Ruby-crowned Kinglet (*Regulus
 calendula*), 90; prehensile
 feet of the Crow (*Corvus frugivo-
 rus*), 92; a Woodpecker destroy-
 ing Cocoons, 93; Are trinomials
 necessary? 101, 200; New Brun-
 swick winter notes, 294; the
 Crow (*Corvus frugivorus*) as a
 fisherman, 391.
 Charadrius dominicus, 179, 248.
 fulvus, 78.
 virginicus, 313.
 Chat, Yellow-breasted, 67, 216, 389.
 Chauna chavaria, 272.
 Chen cærulescens, 240.
 hyperboreus, 34, 240, 386.
 Chickadee, Black-capped, 6, 8, 135,
 157, 244.
 Hudsonian, 185.
 Chimango, 111.
 Chloëphaga magellanica, 273.
 Chloris, 168.
 Chlorostilbon splendidus, 22.
 Chondestes grammacus, 121.
 Chordiles popetue, 246.
 virginianus, 24.
 Chorlito, 313.
 Chorlo, 313, 315.
 Chrysomitris, 361.
 pinus, 160, 176, 177, 328.
 Chrysoptilus cristatus, 25.
 Ciceronia pusilla, 81.
 Ciconia maguari, 271.
 Ciguëña, 271.
 Cinclus aquaticus, 345.
 Circus cinereus, 30.
 hudsonius, 19, 122, 185, 247.
 maculosus, 30.
 Cistothorus stellaris, 59, 139, 244.
 Cisne, 271.
 Clangula albeola, 249.
 glacium americana, 177, 249.
 Clark, J. N., nesting of the Little
 Black Rail in Connecticut, 393.
 Coccyzus cinereus, 28.
 erythrophthalmus, 178, 179.
 melanocoryphus, 28.
 pumilus, 28.
 Colaptes agricola, 25.
 auratus, 246.
 Collins, J. W., notes on certain
 Laridæ and Procellariidæ of the
 New England Coast, 236; notice
 of his paper on Sea Birds used
 as fish-bait by fishermen, 380.

- Columba, 140.
 maculosa, 274.
 picazura, 274.
 Columba picui, 274.
 Colymbus adamsi, 81.
 arcticus, 207.
 glacialis, 115.
 torquatus, 115, 185, 249.
 Compsothlypis americana, 169, 170, 212.
 gutturalis, 169.
 insularis, 169, 170.
 nigrilora, 169, 170.
 pitiayumi, 170.
 pitiayumi inornata, 170.
 Condor, California, 81.
 Contopus borealis, 15.
 Conurus patagonus, 28.
 Cooke, W. W., bird nomenclature of the Chippewa Indians, 242; the distribution and migration of *Zonotrichia querulea*, 332.
 Coot, American, 248.
 Cormorant, 189, 196.
 Double-crested, 249.
 Pallas's, 82.
 Violet-green, 163.
 Corvus corax carinivorus, 160, 245.
 cryptoleucus, 100.
 frugivorus, 115, 160, 245, 391.
 Cory, C. B., descriptions of several New Birds from Santo Domingo, 1, pl. i; notice of his 'Beautiful and Curious Birds,' 81; *Thryothorus ludovicianus* in Massachusetts, 91; *Bernicla brenta nigricans* in Massachusetts, 96; notice of his 'Birds of Haiti and San Domingo,' 285; the generic name *Ligea*, 290.
 Corythus enucleator, 115.
 Cosmonetta histrionica, 115.
 Coturniculus passerinus, 330.
 Coues, E., ornithophilologicalities, 49, 140; notice of his 'A hearing of Bird's Ears,' 182; trinomials are necessary, 197; notice of his 'Key to North American Birds,' second edition, 282; egg of Cowbird in nest of the Carolina Dove, 293; on some new terms recommended for use in Zoölogical Nomenclature, 320; Strickland as an advocate of 'Linnaeus at '58,' 400.
 Coues, E., and D. W. Prentiss, notice of their Avifauna Columbiana, 386.
 Cowbird, 186, 193, 245, 293.
 Crane, Carolina, 399.
 Crane, Corn, 399.
 Crane, Alaskan, 80.
 Sand-hill, 85, 248.
 Whooping, 248, 288.
 Creeper, Black-and-white, 6, 87, 190.
 Crex, Brown, 136, 157, 294.
 pratensis, 399.
 Crispin, 27.
 Crossbill, Red, 88, 159, 292, 327.
 White-winged, 159.
 Crow, 6, 92, 160, 188, 189, 245, 294, 391.
 Carrion, 285.
 Clarke's, 16.
 Cuckoo, 28.
 Cucillo, 28.
 Cuervo, 112.
 del agua, 70.
 Cupidonia cupido, 247, 391.
 Curlew, Bristle-billed, 79.
 Eskimo, 393.
 Cyanocephalus cyanocephalus, 230.
 Cyanocitta cristata, 160, 245.
 Cyanospiza parellina, 88.
 Cygnus americanus, 89, 115.
 columbianus, 115.
 musicus, 173.
 nigricollis, 273.
 Cymochorea leucorrhoea, 98.
 Cypseloides niger borealis, 207.
 DAFILA acuta, 248.
 bahamensis, 274.
 spinicauda, 274.
 Dalgleish, J. J., occurrence of the Royal Tern (*Sterna regia*) at Tangiers in Morocco, 97; a newly-discovered breeding-place of Leach's Petrel (*Cymochorea leucorrhoea*) in Scotland, 98.
 Davison, J. L., Wilson's Petrel in Western New York, 294.
 Dendrocopos purus, 35.
 Dendræca, 47.
 adelaidæ delicata, 83.
 æstiva, 175, 176, 212, 244.
 auduboni, 100.
 blackburniæ, 15, 121, 178.
 cærulescens, 15, 177, 179, 213, 244.
 castanea, 213.
 chrysoparia, 121.
 coronata, 121, 176, 177, 192, 213.
 discolor, 176.
 kirtlandi, 389.
 maculosa, 14, 15, 176, 213.
 palmarum hypochrysea, 213.
 pennsylvanica, 213.
 pinus, 14, 176, 178.

- Dendroica striata*, 176, 177, 178, 179, 213.
tigrina, 59, 213.
virens, 178, 213.
Dicromanassa, 143.
Diplopterus galeritus, 27.
Dodo, 81.
Dolichonyx oryzivora, 176, 245.
Dormilon, 24.
Dove, Mourning, 247.
 Sea, 35.
Dovekie, 35.
Drew, F. M., notes on *Lagopus leucurus*, 392.
Drymornis bridgesi, 20.
Duck, Eider, 184, 289.
 King, 184.
 Labrador, 81.
 Pintail, 248.
 Red-head, 185, 188.
 Scaup, 249.
 Wood, 187, 249.
Dutcher, W., bird notes from Long Island, N. Y., 31, 174.
Dytes auritus, 249.
 nigricollis, 249.
 californicus, 18.
EAGLE, Bald, 247.
 Golden, 10, 247.
 Great Northern Sea, 140.
Ectopistes migratorius, 185, 247.
Egret, American, 32.
Eider, Spectacled, 80.
Elanus glaucus, 115.
 forficatus, 247.
Emberiza nivalis, 118.
Empidonax, 42.
 flaviventris, 15, 295.
Engyptila wellsi, 180.
Eremophila alpestris, 86, 160, 179, 245.
 alpestris chrysolæma, 121, 123.
Erismatura dominica, 274.
Erythacus, 117.
Espatula, 272.
Eudocimus albus, 239.
 ruber, 239.
Eudromias modesta, 313.
Euryornhynchus pygmæus, 78, 81.
FALCO columbarius, 399.
 peregrinus, 397.
 peregrinus naevius, 79, 295, 391.
 peregrinus pealii, 79.
 tinnunculus, 398.
Farlouzanne, 167, 168.
Fewkes, J. W., Do Crows carry objects in their claws? 92; Ducks transporting clams, 195.
Finch, Grass, 6, 8, 11, 31, 330.
 Harris's, 332.
 Lincoln's, 31.
 Pine, 160.
 Purple, 6, 13, 63, 67, 288, 327.
Flicker, Yellow-shafted, 246.
Flycatcher, Canada, 9, 11, 13.
 Couch's Tyrant, 93.
 Great-crested, 7, 65, 86, 391.
 Least, 6.
 Olive-sided, 7, 12.
 Townsend's, 286.
 Yellow-billed, 12.
Fox, W. H., *Dendroica coronata* in Southern New Hampshire in Summer, 192.
Frazer, J. T., the Turkey Buzzard in Western New York, 293.
Fringilla borealis, 152.
 cabaret, 153.
 canescens, 153.
 lapponica, 118.
 linaria, 152.
 linaria alnorum, 153.
 linaria β , 154.
 linaria betularum, 152.
 linaria brevirostris, 152.
 linaria β *holböllii*, 153.
 linaria magnirostris, 153.
 linaria var. β , 153.
 minima, 153.
Fulica, 143.
 americana, 248.
 armillata, 277.
 leucoptera, 277.
Fulix affinis, 249.
 collaris, 249.
 marila, 178, 249.
Fulmar, Slender-billed, 80, 233.
Fulmarus glacialis, 233, 238.
 glacialis glupischa, 234.
 glacialis rogersi, 233.
 glacialoides, 233.
GADOW, H., notice of his Catalogue of the Paridæ and Laniidæ, 279.
Galeoscoptes carolinensis, 175, 176, 244.
Gallina del agua, 277.
Gallinago media wilsoni, 248.
 paraguaiæ, 314.
 wilsoni, 187.
Gallinita chica, 276.
 grande, 276, 277.
Gallinula galeata, 277.
Ganso de la sierra, 273.
Garza, 271.
 blanca, 271.
 blanca chica, 271.
Garzetta, 143.
 candidissima, 32.

- Gavilan, 30.
 Gaviota, 316.
 Gaviotila, 316.
 Geothlypis philadelphia, 14, 15.
 trichas, 175, 176, 177, 178, 179,
 215, 389.
 Gelochelidon nilotica, 365.
 Gill, T., notice of his 'Record of
 Scientific Progress for 1881, '84.
 Golden-eye, American, 249.
 Goldfinch, American, 6, 64, 244,
 329.
 Goniaphea ludoviciana, 14.
 Goose, Blue-winged, 240.
 Canada, 80, 188, 248, 309.
 Emperor, 80.
 Snow, 34, 184, 240.
 White-fronted, 248.
 Goshawk, 247.
 Henshaw's, 166.
 Goss, N. S., notes on the breeding
 habits of the American Eared
 Grebes (*Dytes nigricollis californicus*), 18; birds new to the
 Fauna of Kansas and others rare
 in the State, captured at Wallace,
 Oct. 12 to 16, 1883, 100; notes
 on the nesting of the Yellow-
 throated Vireo (*Lanivireo flavi-*
 frons), 124; notes on *Phalacroco-*
 rax violaceus, and on *P. violaceus*
 resplendens, 163; capture of
 Megalestris skua off the coast of
 Cape Cod, Mass., 395; *Brachy-*
 rhamphus hypoleucus off the coast
 of Southern California, 396.
 Grackle, Purple, 245.
 Graculus bairdi, 165.
 Grebe, American Eared, 18.
 Horned, 249.
 Thick-billed, 122, 249.
 Grosbeak, Black-headed, 367.
 Cardinal, 171, 289, 390.
 Evening, 244.
 Pine, 158, 244, 294, 327.
 Rose-breasted, 8, 9, 70, 245,
 367.
 Grouse, Canada, 247.
 Newfoundland Willow, 369.
 Ruffed, 9, 186, 247.
 Grus americana, 248.
 canadensis, 248.
 fraterculus, 80.
 Guillemot, Xantus's, 396.
 Guira pirigua, 26.
 Gull, Ivory, 80.
 Kittiwake, 380.
 Lesser Glaucous-winged, 196.
 Ring-billed, 19.
 Skua, 395.
 Gull, White-winged, 240, 287.
 Gurney, J. H., Jr., hybrid between
 Pediæetes phasianellus and *Cu-*
 pidonia cupido, 391.
 Gybitz, 120.
 Gyrfalcon, 58, 94.
 HABIA, 366, 367.
 ludoviciana, 367.
 melanocephala, 367.
 Hæmatopus ostrilegus, 172.
 Hag, 237.
 Hagdon, 237.
 Haliaëtus albicilla, 140.
 hypoleucus, 82.
 leucocephalus, 247.
 Harelda, 144, 173.
 glacialis, 177.
 Harporhynchus lecontei, 355, 356.
 rufus, 13, 132, 175, 244.
 Hawk, Broad-winged, 95.
 Cooper's, 247.
 Duck, 391.
 Fish, 86, 247.
 Krider's, 100.
 Marsh, 19, 122, 247.
 Red-shouldered, 186.
 Red-tailed, 123, 247.
 Rough-legged, 247.
 Sharp-shinned, 247.
 Sparrow, 247.
 Swainson's, 247.
 Swallow-tailed, 189.
 Hazard, R. G., 2d, the occurrence
 of the Golden Swamp Warbler
 (*Protonotaria citrea*) in Rhode
 Island, 290; a lay view of orni-
 thophilologicalities, 300.
 Helionaster furcifer, 21.
 Helmintherus, 49.
 Helminthophaga chrysoptera, 186.
 leucobronchialis, 91.
 Helminthophila celata, 121.
 peregrinus, 212.
 pinus, 210, 397.
 ruficapilla, 14, 211.
 Helminthotherus vermivorus, 210.
 Hemiprocne zonaris, 24.
 Helmitherus, 49.
 Hendrickson, W. F., capture of the
 Summer Redbird on Long Island,
 290.
 Henshaw, H. W., description of a
 new Song Sparrow from the
 southern border of the United
 States, 223; on a new Gull from
 Alaska, 250; on the Shore Larks
 of the United States and adjacent
 territory, 254; Indian bird names,
 400.
 Herodias alba egretta, 32.

- Herodias egretta, 59.
 Heron, Great Blue, 247.
 Great White, 59.
 Night, 96, 122.
 Snowy, 32.
 Hesperiphona vespertina, 244.
 Heteractitis incanus, 236.
 Heteroscelus incanus, 79.
 Heterospizias meridionalis, 109.
 Hiaticula, 141.
 Hierofalco, 58.
 gyrfalco, 79.
 gyrfalco sacer, 79, 94.
 Hirundo bicolor, 325.
 erythrogastra, 325, 368.
 saturatus, 82.
 sclateri, 2, 285.
 Histrionicus minutus, 115.
 Holtherhoff, G., *Calamospiza bicolor*
 in Southern California, 293;
 occurrence of the Least Tern at
 San Diego, Cal., 294; Eskimo
 Curlew at San Diego, Cal., 393.
 Hornemanni, 57.
 Hummingbird, Black-chinned, 123.
 Ruby-throated, 246.
 Hydranassa, 143.
 Hydrochelidon lariformis, 59.
 Hydropsalis furcifera, 24.
 Hylocharis sapphirina, 22.
 Hylocichla aliciae, 178, 179.
 fuscescens, 178, 179, 243.
 mustelina, 284.
 unalascae pallasi, 121, 176, 178.
 ustulata swainsoni, 176, 178,
 179, 243.
 Hylotomus pileatus, 246.
 Ibis, Sacred, 81.
 Scarlet, 239.
 White, 239.
 Wood, 297.
 Icteria virens, 216, 389.
 Icterus galbula, 245.
 spurius, 390.
 Ictinia, 58.
 subcærulea, 115, 122.
 Iliacus, 47, 56.
 JAEGER, Arctic, 237.
 Jay, Blue, 85, 86, 160, 245.
 Canada, 160, 245.
 Piñon, 230.
 Jeffries, J. A., notice of his paper on
 the 'Epidermal System of
 Birds,' 182; the occipital style of
 the Cormorant, 196.
 Jencks, F. T., another Gyrfalcon in
 Rhode Island, 94; Can we not
 have a simpler system of nomen-
 clature? 297.
- Jouy, P. L., notice of his paper on
 birds collected in Japan, 387.
 Junco bairdi, 284.
 hiemalis, 14, 15, 177, 178, 179,
 245.
 hiemalis connectens, 283.
 oregonus, 256.
 vulcani, 84.
 KIEBITZ, 120.
 Killdeer, 248.
 Kingbird, 6, 8, 87, 122, 245.
 Kingfisher, 246.
 Kinglet, Golden-crowned, 134, 294.
 Ruby-crowned, 90, 134.
 Kite, Everglade, 95.
 Swallow-tailed, 247, 288, 289.
 Kiwi, 81.
 Knowlton, F. H., the Orchard
 Oriole (*Icterus spurius*) in West-
 ern Vermont, 390.
 Knot, 307.
 Kybitz, 120.
 Kywitz, 120.
 LAGOPUS alba, 79, 82, 225.
 alba alleni, 369.
 alpinus, 80, 225, 227.
 leucurus, 392.
 muta, 226.
 mutus atkhensis, 79.
 rupestris, 79, 225.
 rupestris nelsoni, 226.
 rupestris occidentalis, 79, 225.
 rupestris atkhensis, 226.
 ridgwayi, 226.
 subalpinus, 227.
 Langille, J. H., Bicknell's Thrush,
 268.
 Lanius borealis, 158, 185, 186, 244,
 291, 344.
 bucephalus, 292.
 cristatus, 77, 81, 291.
 ludovicianus, 186, 193.
 major, 292.
 Lanivireo flavifrons, 124, 176.
 solitarius, 122.
 Laopteryx priscus, 84.
 Lark, Desert Horned, 264.
 Horned, 87, 123, 254, 285.
 Louisiana Horned, 167, 168.
 Meadow, 6, 245.
 Prairie Horned, 264.
 Red, 167, 168.
 Ruddy Horned, 267.
 Shore, 89, 160, 189, 245, 254.
 Sprague's, 286.
 Streaked Horned, 267.
 Texan Horned, 266.
 Tit, 89, 186. (See also Tit-
 lark.)

- Larus affinis*, 80, 359.
 argentatus, 232.
 cachinnans, 80, 232.
 chalcopterus, 252.
 cirrhocephalus, 316.
 delawarensis, 19.
 dominicanus, 316.
 fuscescens, 232.
 glaucescens, 287, 360.
 kumlieni, 196, 251.
 leucopterus, 240, 287, 360.
 marinus, 231, 232.
 nelsoni, 250.
 pelagicus, 232.
 schistisagus, 231.
 Lawrence, G. N., characters of a new species of Pigeon of the genus *Engyptila*, from the Island of Grenada, W. I., 180; on the occurrence of the White-winged Gull (*Larus leucopterus* Faber) in the State of New York, 240; notice of his paper on new American Birds, 387.
 Lechuza, 30.
 Lechuzon, 29.
 Lepidocolaptes atripes, 21.
 Leptoptila chaltauchenia, 275.
 Leucibis alba, 236.
 rubra, 236.
 Leuconerpes candidus, 25.
 Ligea, 1, 236.
 palustris, 1, 285.
 Limosa lapponica novæ-zealandiæ, 79.
 uropygialis, 79.
 Linaria, 145, 146.
 agrorum, 152.
 alnorum, 153.
 alnorum magnirostris, 153.
 betularum, 152.
 borealis, 152.
 brunnescens, 153.
 canescens, 153.
 lanceolata, 153.
 longirostris, 153.
 minor, 154.
 rufescens, 154.
 Linnet, Pine, 328.
 Linota canescens, 153.
 exilipes, 153.
 hornemannii, 152.
 rufescens, 154.
 Lintner, G. A., Duck Hawks breeding in the Helderberg Mts., N. Y., 391.
 Lobipes hyperboreus, 33.
 tricolor, 367.
 Loomis, L. M., *Xanthocephalus icterocephalus* in Chester County, South Carolina, 293.
 Loon, 186, 249.
 Lophodytes cucullatus, 249.
 Lophophanes bicolor, 134.
 inornatus cinerascens, 284.
 Lophortyx californicus, 355.
 gambeli, 355.
 Lorito, 28.
 Loro, 28.
 Loxia curvirostra, 88.
 curvirostra americana, 159, 292, 327.
 leucoptera, 159.
 rubra, 171.
 MACHETES pugnax, 186.
 Macrorhamphus griseus scolopaceus, 32.
 Magpie, 245, 286.
 Mallard, 122, 188, 248.
 Black, 248.
 Marbleheader, 238.
 Mareca americana, 394.
 penelope, 80.
 sibilatrix, 274.
 Marling-spike, 237.
 Martin pescador, 26.
 Purple, 6, 244.
 Martinete, 318.
 Mbatitú, 314.
 Mbatitui, 314.
 McCormick, L. M., 'Avifauna Columbiana'—a Protest, 396.
 McIlwraith, T., the Yellow-breasted Chat and Summer Redbird in Canada, 389; Pelicans on the move, 395.
 Melanerpes, 41.
 erythrocephalus, 189, 246.
 formicivorus, 87.
 Meleagris gallopavo americana, 247.
 Melospiza fasciata, 176, 177, 178, 245.
 fasciata fallax, 223.
 fasciata montana, 224.
 lincolni, 31, 121, 175, 295, 397.
 palustris, 14, 176, 295.
 Megalestris skua, 395.
 Mergulus alle, 115.
 Mergus merganser americanus, 249.
 serrator, 177, 249.
 Merle d'eau, 120.
 Merriam, A. C., the Coues Lexicon of North American Birds, 36.
 Merriam, C. H., third addendum to the Preliminary List of the Birds ascertained to occur in the Adirondack Region, Northeastern

- New York, 58; Bird Migration, 71; a plea for the metric system in ornithology, 203; *Vireo philadelphicus* in Northern New York, 291; second addendum to List of Birds ascertained to occur within ten miles from Point de Monts, Province of Quebec, Canada; based chiefly upon the notes of Napoleon A. Comeau, 295; breeding of *Passerculus princeps* on Sable Island, 390.
- Merula confinis, 285.
 migratoria, 161, 178, 244.
 migratoria propinqua, 100.
- Metopiana peposaca, 274.
- Microligea palustris, 290.
- Micropus affinis, 230.
 apus, 230.
 melanoleucus, 230.
 melba, 230.
 pallidus, 230.
 saxatilis, 230.
 unicolor, 230.
- Milvulus tyrannus, 207.
- Mimocichla ardesiaca, 285.
- Mimus carolinensis, 132.
 polyglottus, 131, 192.
- Minor ornithological publications, notices of, 85, 185, 284.
- Mniotilta varia, 176, 190, 210.
- Mockingbird, 69, 131, 192, 286.
- Molothrus, 45.
 ater, 125, 186, 245.
- Morden, J. A., Cowbirds in a Black-and-white Creeper's Nest, 193.
- Morden, J. A., and W. E. Saunders, notice of their 'List of the Birds of Western Ontario,' 85.
- Motacilla, 44, 56.
 amurensis, 387.
 blakistoni, 387.
 eques, 170.
 ludoviciana, 170.
 ocularis, 77, 81, 387.
 yarrelli, 398.
- Myiadestes, 42, 56.
 leucotis, 182.
 montanus, 285.
 reevei, 182.
 townsendi, 91, 100.
- Myiarchus crinitus, 391.
- Myiodioctes canadensis, 15, 178, 217.
 mitratus, 216.
 pusillus, 217.
- NELSON, E. W., notice of his 'Birds of Bering Sea and the Arctic Ocean,' 76; brief diagnoses of two new races of North American birds, 165; the Ornithological Report of the 'Cruise of the Corwin,' 202; the breeding habits of the Pectoral Sandpiper (*Actodromas maculatus*), 218.
- Neocorys spraguei, 121.
- Nettion carolinensis, 248.
- Nighthawk, 246.
- Noddy, 238.
- Nothura maculosa, 317.
- Numenius arquata, 142.
 borealis, 316, 393.
 femorialis, 79.
 taitensis, 79.
- Nuthatch, Red-bellied, 13, 136, 157, 244, 294.
 White-bellied, 135, 157, 244.
- Nyctale acadica, 94, 397.
 tengmalmi, 79.
 tengmalmi richardsoni, 79.
- Nyctea scandiaca, 246.
- Nyctiardea grisea naevia, 96, 122.
- Nycticorax gardeni, 271.
- OCEANITES oceanica, 294, 380.
- Oedemia americana, 178.
- Oedinemus dominicensis, 4.
- Estrelata fisheri, 83, 108.
 gularis, 83.
- Oil-bird, 238.
- Oporornis agilis, 179, 192.
- Oreortyx picta plumifera, 355.
- Oreothlypis gutturalis, 169.
 superciliosa, 169.
- Oriole, 57.
 Baltimore, 6, 8, 65, 245.
 Orchard, 67, 68, 390.
- Ortyx, 141.
 virginianus, 6, 246.
- Ostrælegus, 142.
- Ostrilegus, 142.
- Otocorys alpestris, 257, 263.
 alpestris arenicola, 259, 263, 265.
 alpestris chrysolaema, 260, 263.
 alpestris geraudi, 260, 263, 266.
 alpestris leucolæma, 258, 263, 264.
 alpestris praticola, 258, 263, 264.
 alpestris rubeus, 260, 263, 266.
 alpestris strigata, 261, 264, 267.
 occidentalis, 259.
- Otomela cristata, 292.
- Ovenbird, 6, 8, 9, 13, 213.
- Owl, American Hawk, 363.
 Barn, 246.

- Owl, Barred, 88, 246.
 Burrowing, 286.
 European Hawk, 363.
 Great Gray, 246, 286.
 Great Horned, 194, 246.
 Hawk, 246.
 Little Screech, 246.
 Long-eared, 246.
 Saw-whet, 94.
 Short-eared, 246.
 Snowy, 186, 246.
 White, 185.
Oxyechus vociferus, 248.
PAGOPHILA eburnea, 80.
 Pallasia, 228.
 Paloma, 275.
 del monte, 274.
 Palomita, 274.
 Pandion haliaëtus carolinensis, 79.
 Parisorum, 57.
 Park, A. F., the Lesser Glaucous-winged Gull in New York, 196.
 Parra jacana, 277.
Parula americana, 170, 175, 176, 178, 179.
 brasiliانا, 170.
 inornata, 170.
 insularis, 170.
 nigrilora, 170.
 pitiayumi, 170.
 pitiayumi inornata, 170.
 pitiayumi insularis, 170.
 pitiayumi nigrilora, 170.
Parus americanus, 170.
 atricapillus, 135, 157, 244, 397.
 carolinensis, 397.
 cinctus, 77.
 cinctus griseus, 78.
 hudsonius, 15, 88, 157, 185.
 hudsonius evura, 78, 283.
Passerculus anthinus, 106.
 guttatus, 83.
 princeps, 31, 292, 390.
 rostratus, 83.
 sanctorum, 83.
 sandwichensis savanna, 176, 177, 178, 179, 329.
 Passer, 57.
 domesticus, 87, 159, 208, 295.
Passerella iliaca, 177, 178, 295.
Passerina cyanea, 295.
 Patito, 273, 274.
 Pato, 273.
 Pavo del monte, 274.
Pediæcetes phasianellus, 391.
 Pedincho, 26.
 Pelican, White, 395.
Pelidna subarquata, 32.
Penelope obscura, 275.
 Perdiz, 317.
 grande, 317.
Perisoreus, 57.
 canadensis, 15, 160, 245.
Perspicillata, 144, 173.
 Petrel, Leach's, 98.
 Wilson's, 294.
Petrochelidon pyrrhonota, 368.
 Pewee, Bridge, 6.
 Say's, 122.
 Wood, 6, 8.
Phalacrocorax, 144.
 bicristatus, 189.
 brasilianus, 270.
 dilophus, 249.
 perspicillatus, 82, 144.
 violaceus, 163.
 violaceus resplendens, 163.
 Phalarope, Northern, 33.
 Red, 33.
 Wilson's, 33.
Phalaropus fulicarius, 33, 142.
 wilsoni, 367.
Philohela minor, 248.
Phœnicopterus ignipalliatu, 272.
Phoneus brachyurus, 292.
Phylacte canagica, 80.
Phyllopneuste borealis, 227.
 eversmanni, 227.
 trochilus, 227.
Phylloscopus borealis, 77.
Pica caudata, 115.
 rustica, 115.
 rustica hudsonica, 245.
 Picaflor, 22.
 mayor, 21.
Picicorvus columbianus, 17.
Picoïdes arcticus, 161, 246, 288.
 trydactylus alascensis, 165.
 trydactylus americanus, 161, 295.
Picus cactorum, 25.
 mixtus, 25.
 nuttalli, 207.
 pubescens, 122, 161, 246.
 villosus, 161, 246.
 Pigeon, Passenger, 185, 186, 247.
 Pilincho, 26.
 Pipit, American, 186.
Pinicola enucleator, 158, 244, 327.
 enucleator flammula, 149.
 flammula, 149.
Pipilo erythrophthalmus, 14, 175, 245.
Platalea ajaja, 272.
Plectrophanes, 118.
 nivalis, 159, 177, 178, 245.
Plectrophenax, 119.
Plegadis falcinellus, 272.

- Plotus anHINGA*, 121.
Plover, Golden, 248.
 Killdeer, 248.
 Mongolian, 78.
 Mountain, 183.
Podager nacunda, 24.
Podasocys montanus, 121, 183.
Podiceps holboellii, 177.
 rollandi, 317.
Podilymbus podiceps, 122, 249.
Poliophtila cærulea, 83.
 cærulea obscura, 83.
Polysticta stelleri, 80.
Poæcetes gramineus, 31, 330.
Porzana carolina, 97, 176, 248.
 jamaicensis, 393, 397.
 maruetta, 143, 173.
 noveboracensis, 397.
 Prentiss, D. W., see Coues, E.
Priocella glacialisoides, 233.
 tenuirostris, 80, 233.
Procellaria curilica, 234.
 pacifica, 234.
 tenuirostris, 233, 234.
Progne concolor, 368.
 elegans, 368.
 furcata, 368.
 purpurea, 368.
 subis, 244, 368.
Protonotaria citrea, 290.
Psaltiriparus grindæ, 284.
Pseudotantalus, 83.
Ptarmigan, Alpine, 227.
 Atkhan, 79, 225.
 Rock, 79, 225.
 Subalpine, 227.
Pterocorys, 228.
 Publications received, lists of, 89, 190, 289.
Puffinus fuliginosus, 237, 380.
 major, 237, 380, 381.
 tenuirostris, 108, 223.
Pyrranga æstiva, 290, 390.
 rubra, 14, 178, 179, 244, 326.
Pyrrhula, 48, 56.
 cassini, 77.
Pyrrhuloxia sinuata, 122.
 QUAIL, 186, 188, 247, 287.
 Migratory, 186.
Querquedula, 144.
 brasiliensis, 273.
 cyanoptera, 273.
 discors, 248.
 flavirostris, 273.
 versicolor, 274.
Quiscalus, 57.
 æneus, 86.
 major, 185.
 purpureus, 245.
 RAIL, Little Black, 393.
 Rail, Red-breasted, 97.
 Sora, 248.
 Virginian, 399.
Rallus antarcticus, 276.
 elegans, 97.
 longirostris crepitans, 175.
 maculatus, 276.
 rythrhynchus, 276.
 virginianus, 178, 399.
Raven, 10, 160, 245.
 White-necked, 100.
Redbird, Summer, 390.
Redhead, 249.
Redpoll, American Mealy, 159.
 Lesser, 159.
Redstart, 6, 9, 11, 69, 217.
Regulus calendula, 90, 134, 185.
 satrapa, 134, 177.
Rhea americana, 318.
Rhodostethia rosea, 80.
Rhyacophilus solitarius, 315.
Rhynchæa semicollaris, 314.
Rhynchotus rufescens, 317.
Rhynchophanes maccowni, 121.
 Ridgway, R., notice of his paper on supposed undescribed birds from the Commander Islands, etc., 82; do. on a new Warbler from the Island of Santa Lucia, W. I., 83; do. on a new Plover from Chili, 83; do. on a new Petrel from Alaska, 83; do. on the genus *Tantalus*, 83; do. on a collection of Birds from Costa Rica, 84; note on *Zenaidura yucatanensis* Lawr., 96; note on *Phalacrocorax violaceus resplendens*, 165; remarks upon the close relationship between the White and Scarlet Ibises (*Eudocimus albus* and *E. ruber*), 239; note on *Astur atricapillus striatulus*, 252; on the possible specific identity of *Buteo cooperi* Cass. with *B. harlani* (Aud.), 253; notice of his paper on new birds from Lower California, 284; probable breeding of the Red Crossbill (*Loxia curvirostra americana*) in Central Maryland, 292; probable breeding-place of *Passerculus princeps*, 292; notice of his paper on rare species of Neotropical birds, 386; do. on the Pied Wagtails of Eastern Asia, 387; another Kirtland's Warbler in Michigan, 389.
 Rissa, 144.
 brevirostris, 80, 358.

- Rissa tridactyla*, 380.
tridactyla kotzebuii, 80.
 Robin, 6, 8, 9, 70, 87, 90, 126, 160, 188, 189, 244.
 Western, 100, 123.
Rostrhamus sociabilis plumbeus, 95.
Rupornis ridgwayi, 4.
 SAGE, J. H., another example of *Helminthophaga leucobronchialis* from Connecticut, 91.
Salpinctes obsoletus, 121.
 Sandpiper, Curlew, 32.
 Pectoral, 218, 248.
 Sharp-tailed, 78.
 Saunders, W. E., see Morden, J. A.
 Savanna, 57.
 Sayiornis, 38.
Sayornis fuscus, 179.
 sayi, 122.
Saxicola œnanthe, 295, 378, 398.
Scolopax, 142.
 rusticula, 173.
Scops asio, 246.
 brasiliensis, 29.
 giu, 340.
 Scoter, Velvet, 80, 184.
 Scott, W. L., the winter *Passeres* and *Picariæ* of Ottawa, 156; the generic name *Troglodytes*, 400.
 Sea-hen, 238.
 Sennett, G. B., nest and eggs of Couch's Tyrant Flycatcher (*Tyrannus melancholicus couchi*), 93;
 Black-throated Auk (*Synthliboramphus antiquus* in Wisconsin, 98.
 Seton, Ernest E. T., nest and habits of the Connecticut Warbler (*Oporornis agilis*), 192.
Setophaga ruticilla, 175, 176, 217.
 Shalow, H., notes on '*Lanius cristatus*' and '*L. borealis*' of Nelson's 'Birds of Bering Sea and the Arctic Ocean,' 291.
 Sharpe, R. B., notice of his Catalogue of the *Timeliidæ*, 278; a note on the genus *Progne*, 367.
Shearwater, Greater, 237, 380, 381.
 Slender-billed, 234.
Sheldrake, American, 249.
 Hooded, 249.
 Red-headed, 249.
 Shoveller, 248.
 Shrike, Great Northern, 158, 244, 324.
 Loggerhead, 186, 193, 291.
 Shufeldt, R. W., notice of his paper on the Osteology of *Podiceps montanus*, 183.
Sialia arctica, 121.
 mexicana, 121.
 sialis, 133, 244.
Sitta europæa, 343.
 canadensis, 15, 89, 136, 157, 244.
 carolinensis, 135, 157, 244.
 carolinensis aculeata, 281.
 villosa, 281.
Siurus auricapillus, 13, 175, 176, 178, 179, 213, 295.
 motacilla, 121, 215.
 nævus, 175, 176, 178, 214, 286.
 nævus notabilis, 83.
Skua, Buffon's, 97.
 Great, 238.
 Slade, E., a new element of diagnosis, 402.
 Snipe, Red-bellied, 32.
 Wilson's, 187, 248.
 Snowbird, Black, 9, 11, 13, 245.
 Eastern, 86.
 Solitaire, Townsend's, 100.
 Sparrow, Baird's, 286.
 Chipping, 122.
 English, 85, 87, 159, 188, 189, 287.
 Field, 6, 8, 70.
 Henslow's, 7.
 House, 87, 208, 373.
 Intermediate White-crowned, 100.
 Ipswich, 31, 292.
 Leconte's, 286.
 Mountain Song, 224.
 Savanna, 6, 11, 329.
 Song, 6, 8, 65, 70, 245.
 Swamp, 6, 7, 67.
 Titlark, 106.
 White-crowned, 330.
 White-throated, 11, 12, 122, 331.
 Yellow-winged, 7, 330.
Spatula platalea, 274.
Speotyto cunicularia, 30.
Sphyrapicus varius, 15, 179, 246, 295.
 varius nuchalis, 100.
Spinus lawrencii, 362.
 notatus, 362.
 pinus, 362.
 psaltria, 362.
 psaltria arizonæ, 362.
 psaltria mexicanus, 362.
 tristis, 362.
Spiza townsendi, 191.
Spizella breweri, 121.
 domestica, 175.
 montana, 177.

- Spizella monticola*, 295.
 socialis, 122.
 Starling, Red-winged, 85.
 Stearns, W. A., notice of his 'Notes on the Natural History of Labrador,' 284.
Steatornis caripensis, 307.
Steganopus wilsoni, 33.
 tricolor, 367.
 Stejneger, L., *Dendrocopos purus*, a New Species of Woodpecker from Kamtschatka, 35; notice of his 'Contributions to the History of the Commander Islands. No. 1,' 81; on changes in ornithological nomenclature—a reply to critics, 114; notes on the genus *Acanthis*, 145; *Analecta ornithologica*: I, the occurrence of *Turdus aliciae* in the Palearctic Region, 166; II, on the earliest available name of the American Titlark, 167; III, a brief review of the synonymy of the genus *Compsothlypis* (= *Parula*), 168; IV, on the earliest available name of the Cardinal Grosbeak, 171; V, more 'Ornithophilologicalities,' 172; notice of his 'Systematic Arrangement of the Turdidae,' 181; *Analecta ornithologica* (Second Series): VI, on the Ptarmigans of Nelson's 'Birds of Bering's Sea,' etc., especially those belonging to the group *Attagen* Kaup, 225; VII, on some changes necessary in North American and European ornithological nomenclature, if generic appellations previously applied in botany be not rejected, 228; VIII, *Larus schistisagus*, a new species of Gull from the North Pacific, 231; IX, *Priocella tenuirostris* (Aud.) not a bird of Bering Sea or the Arctic Ocean, 233; X, on old and new generic names, 234; *Analecta Ornithologica* (Third Series): XI, note on Arctic *Lari*, 358; XII, *Chrysomitris* or *Spinus*? 360; XIII, on the systematic name of the American Hawk Owl, 362; XIV, on *Sterna nilotica* of Hasselquist, 364; XV, *Habia* against *Zamelodia*, 366; XVI, on the oldest available name of Wilson's Phalarope, 367; a new subspecies of Willow Grouse from Newfoundland, 369; notice of his paper on 'Trinomials in American Ornithology,' 381.
Stelgidopteryx serripennis, 121.
 Stephens, F., collecting in the Colorado Desert—Leconte's Thrasher, 353.
Stercorarius buffoni, 97, 237.
 pomarinus, 173.
 pomatorhinus, 173.
 skua, 238.
Sterna, 144.
 anglica, 34, 364.
 antillarum, 294.
 cantiaca, 186.
 caspia, 34.
 fuliginosa, 59.
 magnirostris, 316.
 nilotica, 364.
 regia, 97.
 superciliaris, 316.
Sternus cinclus, 120.
Sthenelides melancorypha, 235.
Strepsilas interpres, 173, 399.
Strix canadensis, 362.
 caparoch, 362.
 funerea, 362.
 hudsonica, 363.
Subarquatus, 142.
Sula, 144, 173.
Surnia funerea, 79.
 funerea ulula, 79.
 ulula, 363.
 ulula caparoch, 363.
 Swallow, Barn, 6, 325.
 Cliff, 6.
 White-bellied, 185, 325.
 Swan, Trumpeter, 184.
 Swift, 24.
 Chimney, 13, 161, 188, 246.
Sylvia brasiliana, 170.
 eversmanni, 78, 227.
 pitiavumi, 170.
 plumbea, 170.
 pusilla, 170.
 torquata, 170.
 venusta, 170.
Synthliborhamphus antiquus, 98.
Syrnium cinereum, 286.
 nebulosum, 88.
 TALBOT, D. H., Night Herons and Rails in Dakota, 96.
 Tanager, Scarlet, 8, 9, 13, 64, 190, 244, 326.
 Tantalus, 83.
 loculator, 272.
 Tattler, Wandering, 79.
 Teal, Blue-winged, 248.
 Green-winged, 248.
Telmatodytes palustris, 244.

- Tern, Black, 59.
 Caspian, 34.
 Gull-billed, 34, 365.
 Least, 294.
 Royal, 97.
 Sooty, 59.
 Tero-tero, 278.
 Thalassaëtus pelagicus, 82.
 Thalassoica glacialoides, 80.
 Thamnophilus argentinus, 21.
 cærulescens, 21.
 Theristicus melanopis, 272.
 Thrasher, Brown, 6.
 Leconte's, 353, 355, 356.
 Thrasyaëtus, 41.
 Thrush, Bicknell's 58, 268.
 Brown, 132, 244.
 Golden-crowned, 67.
 Gray-cheeked, 130.
 Hermit, 1, 10, 11, 121, 131.
 Large-billed Water, 215.
 Olive-backed, 12, 13, 129, 243.
 Small-billed Water, 215.
 Wilson's, 6, 8, 9, 11, 129, 243.
 Wood, 6, 8, 9, 11, 128, 190.
 Thryothorus ludovicianus, 91, 137.
 Tijerita-Dormilon, 24.
 Tinnunculus sparverius, 247.
 Tit, Hudson's Bay, 157.
 Titlark, American, 209.
 Titmouse, Tufted, 134.
 Torcaz, 274.
 Totanus cinerascens, 115.
 flavipes, 315.
 glottis, 115.
 melanoleucus, 178, 315, 380.
 Towhee, Arctic, 286.
 Townsend, C. H., notice of his
 'Notes on the Birds of Westmor-
 land County, Penn.,' 184.
 Townsend, C. W., breeding of the
 Mockingbird near Boston, Mass.,
 192.
 Tringa canutus, 295, 307.
 fuscicollis 314.
 maculata, 314.
 merula, 120.
 vanellus, 120.
 Tringoides macularius, 175.
 Trochilus colubris, 123, 246.
 Troglodytes ædon, 137, 244.
 Turdus aliciae, 130, 166.
 aliciae bicknelli, 58, 268.
 fuscescens, 129.
 migratorius, 87, 90, 127.
 mustelinus, 14, 128, 190.
 nanus, 131.
 pallasi, 15.
 pinicola, 181.
 Turdus swainsoni, 15.
 ustulatus swainsoni, 129.
 Turkey, Domestic, 187, 188.
 Wild, 187.
 Turner, M. H., *Vireo philadelphi-*
 cus in the Adirondack Region,
 291.
 Turnstone, 229.
 Black, 229.
 Tyrannus carolinensis, 122, 245.
 intrepidus, 87.
 ULULA cinerea, 79, 246.
 cinerea lapponica, 79.
 flammeata, 362.
 Urinator, 117.
 Urophasianus, 141.
 Urubitinga, 140.
 VANELLUS, 141.
 capella, 120.
 cayennensis, 278.
 Vanneau, 120.
 Vireo, Blue-headed, 332.
 Philadelphia, 291.
 Red-eyed, 6, 8, 9, 11, 65, 217.
 Solitary, 122.
 Warbling, 6, 217.
 White-eyed, 6, 332.
 Yellow-throated, 6, 124, 332.
 Vireo atricapillus, 121.
 flavifrons, 332.
 gilvus, 218.
 noveboracensis, 179, 333.
 olivaceus, 217.
 philadelphicus, 291.
 solitarius, 333.
 Vireosylvia olivacea, 176, 177, 178,
 179.
 Vulture, Black, 286.
 WARBLER, Audubon's, 100.
 Bay-breasted, 213.
 Black - and - white Creeping,
 210.
 Black-and-yellow, 11.
 Blackburnian, 13, 213.
 Black-capped Flycatching,
 217.
 Black-poll, 213, 269.
 Black-throated Blue, 12, 213,
 244.
 Black-throated Green, 9, 11,
 13, 213.
 Blue-winged Yellow, 210, 213.
 Canadian Flycatching, 217.
 Cape May, 59, 213.
 Chestnut-sided, 6, 9, 213.
 Connecticut, 192.
 Golden Swamp, 290.
 Hooded, 216.
 Kirtland's, 389.

- Warbler, Mourning, 9, 12, 13.
 Nashville, 6, 7, 211.
 Prairie, 189.
 Summer, 212.
 Tennessee, 212.
 Worm-eating, 210.
 Yellow, 6, 8.
 Yellow Redpoll, 213.
 Yellow-rumped, 213.
 Yellow-throated, 189.
- Ward, C. W., notes on *Ardea wardi* Ridgw., 161.
- Wasseramsel, 120.
- Water-Thrush. See Thrush.
- Waxwing, Bohemian, 10, 244, 288, 294.
 Cedar, 244.
 Northern, 158.
- Welsh, F. R., odd nesting-site of the Great-crested Flycatcher, 391.
- Wheatear, 378, 398.
- Whippoorwill, 246.
- Widgeon, American, 394.
 European, 80.
- Wife, Old, 184.
- Willard, S. W., the migration of our winter birds, 221.
- Wilsonia bonapartii, 231.
 minuta, 231.
 mitrata, 231.
- Wilsonia pusilla, 231.
- Wintle, E. D., the Loggerhead Shrike (*Lanius ludovicianus*), 193.
- Woodcock, 87, 248.
- Woodpecker, Alaskan Three-toed, 164.
 Banded-backed Three-toed, 161.
 Black-backed Three-toed, 93, 161.
 Downy, 9, 93, 122, 161, 246.
 Golden-winged, 6.
 Hairy, 9, 12, 87, 161, 246.
 Ivory-billed, 286.
 Pileated, 9, 12, 246.
 Red-headed, 189, 246.
 Red-naped, 100.
 Yellow-bellied, 12, 86, 286.
- Wren, Carolina, 137, 286.
 House, 6, 85, 137, 244.
 Long-billed Marsh, 138.
 Short-billed Marsh, 59, 139.
 Winter, 12, 13, 138.
- YELLOW-THROAT, Maryland, 389.
- Young, C. J., notes on certain birds observed on a voyage from Liverpool to Quebec in September, 1883, 398.
- ZAMELODIA, 366.

ERRATA.

- Page 31, line 16, transpose the words *Extent* and *Length*.
- " 32, " 14, for 1881 read 1882.
- " 81, " 38, for Smidt read Smit.
- " 96, last word, for *grævia* read *nævia*.
- " 105, line 35, dele the words 'publication of the.'
- " 106, " 3, for of read on.
- " 185, " 6, add —C. H. M.
- " 208, " 4, for Pickering read Picking.
- " 223, " 33, for collection read Museum.
- " " 34, for Museum read collection.
- " 260, " 40, for 8 read 7.
- " 261, " 20, for 9 read 8.
- " 290, " 25, for mein read mien.
- " 303, " 36, for naturalist read naturalists.
- " 308, " 28, for Blackiston read Blakiston.
- " " 46, for 1877 read 1887.
- " 406, " 9, (first column) for Astrigalinus read Astragalinus.